

Fuzzy bi-ideals in semigroups

by

Nobuaki KUROKI

(Received September 29, 1978)

The concept of a fuzzy set, introduced in Zadeh [5], was applied to the elementary theory of groupoids and groups in Rosenfeld [4] and of semigroups in the author [2]. In the present note we shall give a characterization of a semigroup which is a group, a union of groups and a semilattice of groups in terms of fuzzy bi-ideals.

A *fuzzy set* in a semigroup S is a function δ from S into the unit interval $[0, 1]$. A fuzzy set δ in S is called a *fuzzy subsemigroup* of S if

$$\delta(xy) \geq \min \{ \delta(x), \delta(y) \}$$

for all $x, y \in S$. A fuzzy subsemigroup δ of S is called a *fuzzy bi-ideal* of S if

$$\delta(xyz) \geq \min \{ \delta(x), \delta(z) \}$$

for all x, y and $z \in S$. A subsemigroup A of a semigroup S is called a *bi-ideal* of S if $ASA \subseteq A$. The following theorem shows that the concept of a fuzzy bi-ideal in a semigroup is an extended one of a bi-ideal.

THEOREM 1. *For a non-empty subset A of a semigroup S the following conditions are equivalent.*

- (1) *A is a bi-ideal of S .*
- (2) *The characteristic function δ_A of A is a fuzzy bi-ideal of S .*

Proof. Assume that (1) holds. Let x, y and z be any elements of S . If $x \in A$ and $z \in A$, then we have

$$\delta_A(x) = \delta_A(z) = 1.$$

Since A is a bi-ideal of S ,

$$xyz \in ASA \subseteq A.$$

Then we have

$$\delta_A(xyz) = 1 = \min \{ \delta_A(x), \delta_A(z) \}.$$

If $x \notin A$ or $z \notin A$, then

$$\delta_A(x) = 0 \quad \text{or} \quad \delta_A(z) = 0.$$

Then we have

$$\delta_A(xyz) \geq 0 = \min \{ \delta_A(x), \delta_A(z) \} .$$

Then it follows from Proposition 2.2 of [4] that the characteristic function δ_A is, if A is a subsemigroup of S , a fuzzy subsemigroup of S . Thus we obtain that δ_A is a fuzzy bi-ideal of S , and that (1) implies (2). Conversely, assume that (2) holds. Let $xyz(x, z \in A, y \in S)$ be any element of ASA . Since $x, z \in A$, we have

$$\delta_A(x) = \delta_A(z) = 1 .$$

Then, since δ_A is a fuzzy bi-ideal of S , we have

$$\delta_A(xyz) \geq \min \{ \delta_A(x), \delta_A(z) \} = 1 .$$

This implies that

$$\delta_A(xyz) = 1 ,$$

and so $xyz \in A$. Thus we obtain that $ASA \subseteq A$. It follows from Proposition 2.2 of [4] that the non-empty subset A of S is, if δ_A is a fuzzy subsemigroup of S , a subsemigroup of S . Thus we obtain that A is a bi-ideal of S , and that (2) implies (1).

THEOREM 2. *For a semigroup S the following conditions are equivalent.*

- (1) S is a group.
- (2) Every fuzzy bi-ideal of S is a constant function.

Proof. First assume that S is a group. Let δ be any fuzzy bi-ideal of S , and a any element of S . We denote by e the identity of the group S . Then we have

$$\begin{aligned} \delta(a) &= \delta(eae) \geq \min \{ \delta(e), \delta(e) \} = \delta(e) \\ &= \delta(ee) = \delta \{ (aa^{-1})(a^{-1}a) \} \\ &= \delta \{ a(a^{-1}a^{-1})a \} \geq \min \{ \delta(a), \delta(a) \} \\ &= \delta(a) , \end{aligned}$$

and so we have

$$\delta(a) = \delta(e) .$$

This means that δ is a constant function. Thus (1) implies (2). Conversely, assume that S is not a group. Then it follows from p. 84 of [1] that S contains a proper bi-ideal A of S . Since A is non-empty, the characteristic function δ_A of A is not a constant function. Then, since δ_A is a fuzzy bi-ideal of S by Theorem 1, (2) does not hold. Thus (2) implies (1). This completes the proof.

A semigroup S is called *completely regular* if, for each element a of S , there exists an element x in S such that

$$a = axa \quad \text{and} \quad ax = xa .$$

The following characterization of such a semigroup is due to p. 105 of [6].

LEMMA 3. For a semigroup S the following conditions are equivalent.

- (1) S is completely regular.
- (2) S is a union of groups.
- (3) $a \in a^2Sa^2$ for all $a \in S$.

Now we give another characterization of a completely regular semigroup.

THEOREM 4. For a semigroup S the following conditions are equivalent.

- (1) S is completely regular.
- (2) For every fuzzy bi-ideal δ of S ,

$$\delta(a) = \delta(a^2)$$

holds for all $a \in S$.

Proof. First assume that (1) holds. Let δ be any fuzzy bi-ideal of S and a any element of S . Then there exists an element x in S such that

$$a = a^2xa^2 .$$

Then, since δ is a fuzzy bi-ideal of S , we have

$$\begin{aligned} \delta(a) &= \delta(a^2xa^2) \geq \min \{ \delta(a^2), \delta(a^2) \} \\ &= \delta(a^2) \geq \min \{ \delta(a), \delta(a) \} = \delta(a) , \end{aligned}$$

and so we have

$$\delta(a) = \delta(a^2) .$$

Thus (1) implies (2). Conversely, assume that (2) holds. We denote by $B[x]$ the principal bi-ideal of a semigroup S generated by x in S , that is,

$$B[x] = \{x\} \cup \{x^2\} \cup xSx .$$

Since the characteristic function $\delta_{B[a^2]}$ of the bi-ideal $B[a^2]$ is a fuzzy bi-ideal of S by Theorem 1, and since $a^2 \in B[a^2]$, we have

$$\delta_{B[a^2]}(a) = \delta_{B[a^2]}(a^2) = 1 .$$

This implies that

$$a \in B[a^2] = \{a^2\} \cup \{a^4\} \cup a^2Sa^2 .$$

Then it is easily to see that S is completely regular. Thus (2) implies

(1). This completes the proof.

The following is due to Theorem 1 of [3].

LEMMA 5. *A semigroup S is a semilattice of groups if and only if the set of all bi-ideals of S is a semilattice under the multiplication of subsets.*

THEOREM 6. *For a semigroup S the following conditions are equivalent.*

- (1) *S is a semilattice of groups.*
- (2) *For every fuzzy bi-ideal δ of S ,*

$$\delta(a) = \delta(a^2) \quad \text{and} \quad \delta(ab) = \delta(ba)$$

hold for $a, b \in S$.

Proof. Assume that (1) holds. Then S is a union of groups. Then it follows from Lemma 3 that S is completely regular. And so it follows from Theorem 4 that for every fuzzy bi-ideal δ of S

$$\delta(a) = \delta(a^2)$$

holds for all $a \in S$. Let a and b be any elements of S . Then by Lemma 5 we have

$$\begin{aligned} (ab)^3 &= (aba)(bab) \in B[aba]B[bab] \\ &= B[bab](B[aba])^2 \subseteq B[bab]SB[aba] \\ &\subseteq babSaba \subseteq baSba. \end{aligned}$$

This implies that there exists an element x in S such that

$$(ab)^3 = (ba)x(ba).$$

Then, for any fuzzy bi-ideal δ of S , we have

$$\begin{aligned} \delta(ab) &= \delta\{(ab)^3\} = \delta\{(ba)x(ba)\} \\ &\geq \min\{\delta(ba), \delta(ba)\} = \delta(ba). \end{aligned}$$

Similarly, we can prove that

$$\delta(ba) \geq \delta(ab).$$

Thus we obtain that

$$\delta(ab) = \delta(ba)$$

and that (1) implies (2). Conversely, assume that (2) holds. Then it follows from the first condition and from Theorem 4 that S is completely regular. Then it is easily proved that every bi-ideal of S is globally idempotent. Let A and B be any bi-ideals of S , and ba ($a \in A$, $b \in B$) any element of BA . Then, since by Theorem 1 the characteristic function $\delta_{B[ab]}$ of the bi-ideal $B[ab]$ is a fuzzy bi-ideal of S ,

$$\delta_{B[ab]}(ba) = \delta_{B[ab]}(ab) = 1 .$$

This implies that

$$ba \in B[ab] = \{ab\} \cup \{abab\} \cup abSab .$$

Then

$$(i) \quad ba = ab \in AB, \quad (ii) \quad ba = abab \in ABAB \subseteq AB, \quad \text{or}$$

(iii) Since the product AB of the bi-ideals A and B of S is also a bi-ideal of S , we have

$$ba \in abSab \subseteq (AB)S(AB) \subseteq AB .$$

In any cases we have

$$BA \subseteq AB .$$

Similarly we can see that the converse inclusion holds. Thus we obtain that

$$AB = BA ,$$

and that the set of all bi-ideals of S is a commutative idempotent semigroup. Therefore it follows from Lemma 5 that S is a semilattice of groups, and that (2) implies (1).

COROLLARY 7. *For an idempotent semigroup S the following conditions are equivalent.*

- (1) S is commutative.
- (2) For every fuzzy bi-ideal δ of S ,

$$\delta(ab) = \delta(ba)$$

holds for all $a, b \in S$.

References

- [1] CLIFFORD, A. H. and PRESTON, G. B.; *The Algebraic Theory of Semigroups, Vol. I*, Math. Surveys No. 7, Amer. Math. Soc. Providence, R. I., (1961).
- [2] KUROKI, N.; On fuzzy ideals and fuzzy bi-ideals in semigroups, to appear in the International Journal for Fuzzy Sets and Systems.
- [3] LAJOS, S.; A note on semilattice of groups. *Acta Sci. Math. Szeged*, **33** (1972), 315-317.
- [4] ROSENFELD, A.; Fuzzy groups. *J. Math. Anal. Appl.*, **35** (1971), 512-517.
- [5] ZADEH, L. A.; Fuzzy sets. *Inform. Control*, **8** (1965), 338-353.
- [6] PETRICH, M.; *Introduction to Semigroups*, Columbus, Ohio, (1973).

Department of Mathematics
 General Education
 College of Science and Technology
 Nihon University
 Narashinodai, Funabashi-shi,
 Chiba-ken, Japan